

ADAMSTM *plus*

YARD SPRAY

Premise Control STEP

- ◆ **Kills & repels: fleas, ticks, mosquitoes, ants, crickets & other listed insects**
- ◆ **Treats up to 5,000 sq. ft.**
- ◆ **For use on lawns, trees, shrubs, roses & flowers**
- ◆ **For use around residential (home) areas only**

ACTIVE INGREDIENTS:

Permethrin* 2.5%

OTHER INGREDIENTS: 97.5%

TOTAL: 100.0%

*Cis/trans isomer ratio: Min. 35% (±) cis,
Max. 65% (±) trans

**Kills and Repels
Mosquitoes**



**Up To 4
Weeks Control**

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Back Panel for Additional Precautionary Statements

Net Contents: 1 Qt (32 fl oz)

RM #2008324 06-1904

ADAMSTM *plus* YARD SPRAY



PRECAUTIONARY STATEMENTS - HAZARDS TO HUMANS AND DOMESTIC ANIMALS - CAUTION: May be harmful if swallowed or absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.

EPA Reg. No. 28293-271-270 EPA Est. No. 59144-GA-1



Manufactured For:
Farnam Pet Products
A Division of Farnam Companies Inc.
P. O. Box 34820
Phoenix, AZ 85067-4820



©2006 Farnam Companies, Inc. Made in USA
Product Code: 3006022 RM #: 2008325 06-1904 0 39079106022 6

Pg. 1

FIRST AID (continued)

If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
-------------------	---

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish. Do not apply directly to water. Drifts and runoff from treated areas may be hazardous to aquatic organisms in treated areas. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

STORAGE AND DISPOSAL

STORAGE: Store in a cool, dry area away from heat or open flame. **DISPOSAL: If empty** – Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled** – Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product kills troublesome outdoor insects which destroy lawns, roses, flowers, trees and shrubs: Ants, Bagworms, Aphids, Armyworms, Chinch bugs, Inchworms, Spider mites, Beetles, Fleas, Cicadas, Thrips, Cutworms, Sod webworms, Leafrollers, Whiteflies, Moths.

Pg. 2

Also kills Brown dog ticks, Budworms, Citrus black flies, Cluster flies, Crickets, Earwigs, Elm spanworms, Fall cankerworms, Gnats, Grasshoppers, House flies, Japanese beetles, Leafminers, Lone star ticks, Mealybugs, Millipedes, Mole crickets, Mosquitoes, Scales, Sowbugs (pillbugs), Tent caterpillars, Ticks including Deer tick (which may carry Lyme disease), and many others.

HOSE END SPRAYER: Easy to Use. For Application to Lawns, Outdoor Ornamentals, Outside Surfaces of Buildings, and Other Outdoor Residential Areas. 32 ounces treat up to 5,000 square feet.

Snap off tab on top of sprayer before first use.

1. Make sure knob Control lever is in the **OFF** position. Shake well and attach to hose.
2. Turn water on at faucet, aim nozzle toward lawn and turn knob Control lever to **ON** to begin spraying.
3. When finished, turn Control lever to **OFF** position. Turn faucet to **OFF** position. Discharge residual water pressure by turning **ON** and **OFF** again. Store out of reach of children.

Do not spray animals. People and pets should not be allowed in treated areas until spray has dried.

3 EASY STEPS TO MIX AND APPLY:

- #1:** For lawns, measure the area to be sprayed. Calculate square feet by multiplying length by width. For all other applications, apply as a thorough cover spray.
- #2:** Use a clean sprayer. Carefully measure and mix the amount of concentrate and water as directed.

Pg. 3

#3: Spray as directed. Thorough coverage is important. Flush sprayer with clean water after each use. **Make applications when insects first appear.** Repeat applications as directed. See directions for complete insect list.

Use dilution rates as indicated. 1 fl. oz. = 6 tsp. or 2 Tbsp.; 8 fl. oz. = 1/2 pint or 1 cup. Food utensils such as teaspoons or measuring cups should not be used for food purposes after use with insecticides.

NUISANCE PESTS IN OUTDOOR AREAS			
TREATMENT AREAS	PESTS	REMARKS	USE RATE
Outside surfaces of buildings	Ants (including foraging fire ants), Chiggers, Clover mites, Crickets, Earwigs, Millipedes, Sowbugs (pillbugs), Mosquitoes	To help prevent infestation of buildings, treat the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Also, apply as a residual spray to outside surfaces of building, including porches, screens, window frames, eaves, patios, garages, refuse dumps and other areas where these pests congregate or have been seen. Repeat treatment as needed to maintain effectiveness.	6 2/3 fl. oz. in 1 quart of water, treating 400 linear feet with a 6-inch spray band.
Other Outdoor areas	Ants (including foraging fire ants), Chiggers, Crickets,	For treatment of localized infestations of these insects in areas where there are weeds or bushy non-crop areas,	

Pg. 4

TREATMENT AREAS	PESTS	REMARKS	USE RATE
Other Outdoor areas (continued)	Earwigs, Fleas, Lone star ticks, Ticks, including Deer tick (which may carry Lyme disease)	spray infested areas thoroughly. For Ants , thoroughly wet hills and runways. Repeat application as reinfestation occurs.	6 2/3 fl. oz. in 1 quart of water, treating 400 linear feet with a 6-inch spray band.
	Cluster flies, Flying moths, Gnats, House flies, Mosquitoes	For use only as an aid in reducing annoyance from these insects. Spray outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, in garages and other places where these insects alight or congregate. Repeat as necessary.	
	Tent Caterpillars	Application should be made when caterpillars are young and tents are first noticed. For best results, apply in the late afternoon or evening, when caterpillars have returned to their tents. Wet tents with spray on all sides. A few caterpillars may be seen escaping from the tent, however, they should die within several hours.	

Pg. 5

		HOME LAWN	
TREATMENT AREAS	PESTS	REMARKS	USE RATE
Bent, Bermuda, Bluegrass, Dichondra, Fescue, Irish Moss, Merion, St. Augustine	Armyworms, Brown dog ticks, Chiggers, Chinch bugs, Cutworms, Fleas, Japanese beetle grubs, Mole crickets, Mosquitoes, Sod webworms, Ticks, including Deer tick (which may carry Lyme disease)	Thoroughly wet down grass a few hours before applying. Home lawns should be no taller than 3 inches at time of application. Slightly more water may be used as long as the amount of product per area is as listed below. For heavy infestations, repeat application after 2 weeks.	6 fl. oz. in 10 gallons of water to cover 1,000 sq. ft.
	Ants (including foraging fire ants), Crickets, Grasshoppers		2 fl. oz. in 3 gallons of water to cover 1,000 sq. ft.
	Fire Ants (mound treatment)	Apply 1 gallon of solution as a gentle rain to each fire ant mound using a sprinkler can. Thoroughly wet the mound and surrounding area to a 4 foot diameter. For best results, apply in cool weather, 65° - 80° F, or in early morning or late evening hours. Treat new mounds as they appear. Do not	6 fl. oz. in 1 gallon of water to treat one mound.

Pg. 6

TREATMENT AREAS	PESTS	REMARKS	USE RATE
Bent, Bermuda, Bluegrass, Dichondra, Fescue, Irish Moss, Merion, St. Augustine (continued)		disturb the mound prior to treatment. If possible, treat all mounds in the vicinity, and treat all colonies which may not as yet have constructed a mound. Larger quantities of solution can be prepared for multiple mound treatments. Equipment to deliver the diluted product as a gentle rain at the rate of 1 gallon per mound is still required. Do not use equipment that produces a pressurized spray as pressurized sprays may disturb fire ants and cause migration, reducing product effectiveness.	6 fl. oz. in 1 gallon of water to treat one mound.
TREES & SHRUBS			
TREATMENT AREAS	PESTS	REMARKS	USE RATE
Arizona cypress, Azalea, Birch, Cherry, Non-bearing citrus, Conifers, Elm, English ivy, Euonymus, Weeping fig, Fir, Honeysuckle, Lilac, Mock-orange,	Aphids, Bagworms, Cicadas, Exposed thrips, Fall canker- worms, Inchworms, Leafminers, Leaf- rollers, Mealybugs, Spider mites, Whiteflies	Apply as a thorough spray, wetting leaves and branches to the dripping point. Try to penetrate dense foliage. Spray in the late afternoon or evening when the temperature ranges from 50° to 75° F and when there is little or no wind. Spray at the first sign of insects. Repeat as necessary; use intervals of	1 fl. oz. in 1 gallon of water

Pg. 7

TREATMENT AREAS	PESTS	REMARKS	USE RATE	
Oak, Palm, Philodendron, Pine, Poinsettia, Tulip, Poplar, Rhododen- dron, Taxus Cherry, Honeysuckle Conifers Elm Douglas & Grand firs Non-bearing citrus Oak Pine		4-8 days. Application can be made up to the day of collection.	1 fl. oz. in 1 gallon of water	
	Fall webworms			
	Budworms			
	Elm leaf beetles, Elm spanworms, Fall webworms			
	Tussock moths			
	Citrus black flies			
	Oakworms			
	Pine beetles, Pine moths, Pine needle- miners, Needle scales			
	ROSES & FLOWERS			
	TREATMENT AREAS	PESTS	REMARKS	USE RATE
Ageratum, Aster, Azalea, Begonia, Coleus, Common ninebark and snowberry, Exacum, Gladiolus, Gold bells,	Aphids, Bagworms, Cicadas, Exposed thrips, Fall canker- worms, Fall web- worms, Inchworms, Leafminers,	Apply as a thorough spray, wetting the plants to the dripping point. Try to hit underside of leaves and penetrate dense foliage. Spray in the later afternoon or evening, when the temperature ranges from 50°	1 fl. oz. in 1 gallon of water	

Pg. 8

TREATMENT AREAS	PESTS	REMARKS	USE RATE
Hypoestes, Ivy, Lilac, Marigold, Mock-orange, Nannyberry, Orchid, Pansy, Pea shrub, Petunia, Poinsettia, Rhododendron, Rose, Snapdragon, Zinnia	Leafrollers, Mealybugs, Spider mites, Whiteflies	to 75° F and when there is little or no wind. Spray at the first sign of insects. Repeat as necessary; use intervals of 4-8 days. Application can be made up to the day of collection.	1 fl. oz. in 1 gallon of water

For information or in an emergency, call **1-800-234-2269**.
adamsfleacontrol.com

Notice: Buyer assumes all responsibility for safety and use not in accordance with directions.



Manufactured For:
 Farnam Pet Products
 A Division of Farnam Companies Inc.
 P. O. Box 34820
 Phoenix, AZ 85067-4820

3.5"
Pg. 9
glued base