

SER
EXP

Indications: Recommended for the vaccination of healthy, susceptible cattle and sheep as an aid in the reduction of diseases caused by *Clostridium chauvoei*, *Cl. septicum*, *Cl. haemolyticum*, *Cl. novyi*, *Cl. sordellii*, and *Cl. perfringens* Types C & D. Although *Cl. perfringens* Type B is not a significant problem in the U.S.A., immunity may be provided against the beta and epsilon toxins elaborated by *Cl. perfringens* Type B. This immunity is derived from the combination of Type C (beta) and Type D (epsilon) fractions.



Manufactured by:
Boehringer Ingelheim Vetmedica, Inc.
St. Joseph, MO 64506 U.S.A.
U.S. Veterinary License No. 124
Manufactured for:
Agri Laboratories, Ltd.
St. Joseph, MO 64503 U.S.A.

**CLOSTRIDIUM
CHAUVOEI-SEPTICUM-
HAEMOLYTICUM-
NOVI-SORDELLII-
PERFRINGENS
TYPES C & D
BACTERIN-TOXOID**

CLOSTRIDIAL 8-WAY

250 mL
50 Cattle Doses or 100 Sheep Doses



Composition: Prepared from cultures of the organisms listed. Alum precipitated.

Dosage:

Cattle: Using aseptic technique, inject 5 mL subcutaneously. Repeat in 21 to 28 days and once annually.

Sheep: Using aseptic technique, inject 2.5 mL subcutaneously. Repeat in 21 to 28 days and once annually.

Precautions: Store out of direct sunlight at 35-45°F (2-7°C). Avoid freezing. Shake well before using. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Anaphylactoid reactions may occur. Antidote: Epinephrine.

24214-00

