

# Cetirizine

*Dr. Reedy's*

Hydrochloride Tablets USP, 10 mg  
Antihistamine

Indoor & Outdoor Allergies

Relief of:

- Sneezing
- Runny Nose
- Itchy, watery eyes
- Itchy throat or nose

## Drug Facts

**Active Ingredient (in each tablet)**      **Purpose**

Cetirizine HCl USP, 10 mg...Antihistamine

USES: Temporarily relieves these symptoms due to hay fever or other respiratory allergies:

- Sneezing
- Itchy, watery eyes
- Itching of the nose or throat

DO NOT USE if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

ASK A DOCTOR BEFORE USE IF YOU HAVE liver or kidney disease. Your doctor should determine if you need a different dose.

ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE taking tranquilizers or sedatives.

## WHEN USING THIS PRODUCT

- Drowsiness may occur
- Avoid alcohol, sedatives, and tranquilizers may increase drowsiness
- Be careful when driving a motor vehicle or operating machinery

STOP USE AND ASK A DOCTOR IF an allergic reaction to this product occurs.  
Seek medical help right away.

**NDC 43598-811-15**

**Original Prescription Strength**

**24-hour ALLERGY**

**500 Tablets**

**IF PREGNANT OR BREASTFEEDING:**

- If breast-feeding: not recommended
- If pregnant: ask a health professional before use.

**KEEP OUT OF REACH OF CHILDREN**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions:**

Adults and Children 6 Years and Over	One 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.
Adults 65 Years and Over	Ask a doctor
Children Under 6 Years of Age	Ask a doctor
Consumers with Liver or kidney disease	Ask a doctor

Other information: Store between 20° to 25°C (68°to 77°F)

**Inactive Ingredients:**

Hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, povidone, starch, titanium dioxide

**Questions?** Call 1-888-375-3784