

# Beef Cattle Max

## Weaver Livestock

### Ractopamine Medicated Finishing Cattle Feed

(Weight Gain, Feed Efficiency, and Carcass Leanness) (ractopamine hydrochloride Type B medicated feed)

#### Indications for use:

For increased rate of weight and improved feed efficiency, and for increased rate of weight gain, improved efficiency, and increased carcass leanness in cattle feed in confinement for slaughter during the last 28 to 42 days on feed.

Indications	Appropriate Concentration of Ractopamine in Type C Medicated Feed
Increased Rate of Weight Gain and Improved Feed Efficiency	8.2 to 24.6 g/ton (9ppm to 27ppm)
Increase Rate of Weight Gain, Improved Feed Efficiency and increased Carcass Leanness	9.8 to 24.6 g/ton (11 ppm to 27 ppm)

#### Type B Medicated Feed

Active Drug Ingredient: Ractopamine Hydrochloride .....4,000 g/ton

Ingredients: Rice Hulls, Calcium Carbonate, and Mineral Oil.

The active ingredient in Optaflexx, Ractopamine Hydrochloride is a beta-adrenergic agonist. Individuals with cardiovascular disease should exercise special caution to avoid exposure. Not for use in humans. Keep out of reach of children. The Optaflexx 45 formulation (Type A Medicated Article) poses a low dust potential under usual conditions of handling and mixing. When mixing and handling Optaflexx, use protective clothing, impervious gloves, protective eyewear, and a NIOSH approved dust mask. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse eyes thoroughly with water. If irritation persists, seek medical attention. The material safety data sheet contains more detailed occupational safety information. To report adverse effects, access medical information, or obtain additional product information call, 1-800-428-4441.

#### Guaranteed Analysis:

Crude Protein, not less than .....1.0%  
 Crude Fat, not less than .....1.0%  
 Crude Fiber, not more than .....25.0%  
 Calcium, not less than .....14.0%  
 Calcium, not more than .....17.0%  
 Phosphorus, not less than.....0.1%

### Mixing & Feeding Directions:

Thoroughly mix Bactopamine Type B Medicated Feed into one ton of complete cattle feed according to the table below to obtain the proper concentration in the Type C Medicated Feed. Prepare an intermediate pre-blend of the medicated feed prior to mixing in a complete feed. Thoroughly mix the required amount in a convenient quantity of feed ingredients, then add to the remaining feed ingredients to make a ton of complete feed. [Use only the portion of the table below on your Type B Medicated Feed product label that is applicable to the concentration of Ractopamine in the Type B Medicated Feed which you manufacture.]

Concentration of Ractopamine in Type B Medicated Feed grams/ton	Pounds Type B Medicated Feed to Add per Ton of Type C Medicated Feed	Resulting Ractopamine Concentration in Type C Medicated Feed grams/ton
4000	4.1	8.2
4000	12.3	24.6
4000	4.9	9.8
4000	12.3	24.6

**Caution:** Not for animals intended for breeding.

**Important:** Must be thoroughly mixed into feed before use. Do not feed undiluted.

Withdrawal periods: No withdrawal period is required when used according to labeling.