

HERD HEALTH OPPORTUNITIES:

OPTIONS FOR SPRING CALVING HERDS

Your herd health veterinarian should be consulted to build a customized protocol based on your specific herd requirements.

CLASS/ STAGE OF PRODUCTION	ACTION/VACCINATION	ELANCO PRODUCT OPTIONS	DATE OF ADMINISTRATION AND COMMENTS
Calves: At birth	Colostrum Intake	Calves should be up and nursing 30-60 minutes after birth. If in doubt, tube calf with 2-4 quarts colostrum as soon as possible.	
	Respiratory: Viral	MLV: Titanium® 5 OR Inactivated: Vira Shield® 6	
Calves: 3 Months / Branding	Respiratory: Bacterial	Nuplura® PH. Can also be given at weaning. Consult your herd health veterinarian.	
	7- or 8-way Clostridial	Per herd health veterinarian recommendation.	
	Implant	Component® E-C with Tylan to all steers and non-replacement heifers over 45 days of age.	
	Parasite/Fly Control	Permethrin™ CDS Pour-On OR CyLence® OR Clean Up™ II Pour-On Apply one Corathon® OR CyLence Ultra® insecticide ear tag.	
Calves: Pre-weaning/ Weaning	Respiratory: Viral	MLV: Titanium 5 OR Nuplura PH + 3	
	Respiratory: Bacterial	Nuplura PH. Can also be given at branding. Consult your herd health veterinarian.	
	7- or 8-way Clostridial	Per herd health veterinarian recommendation	
	Parasite/Fly Control**	Cyductin® (moxidectin) Pour-On Clean Up II Pour-On 2-3 weeks later for optimal louse control OR Cyductin® (moxidectin) Injectable Clean Up II Pour-On Insecticide with IGR	
Heifers: Pre-breeding	Respiratory/ Reproductive Vaccine	Inactivated: Vira Shield® 6 VL5 HB OR Vira Shield 6 L5 HB Other Vira Shield combinations are available. Consult your herd health veterinarian.	
	Parasite/Fly Control	Cyductin Pour-On OR Cyductin Injectable Permethrin CDS Pour-On OR CyLence OR Clean Up II Pour-On*	
		Apply two Corathon OR CyLence Ultra insecticide ear tags. Start feeding Rabon® Oral Larvacide in free choice mineral 3-4 weeks before fly season.	

CONTINUED ON NEXT PAGE >>>

For all products: The label contains complete use information, including cautions and warnings. Always read, understand and follow the label and use directions.

Directions for Use: Implants are indicated for increased rate of weight gain; see product labels for full indications. Administer one dose in the ear subcutaneously according to label directions. See label for revaccination interval recommendations.

* If you dewormed in the spring you may not need to deworm in the fall. Talk to your herd health veterinarian regarding a strategic deworming program for your herd.

Labels contain complete use information, including cautions and warnings. Always read, understand and follow the label and use directions.

** Dependent on region.

OPTIONS FOR SPRING CALVING HERDS

Your herd health veterinarian should be consulted to build a customized protocol based on your specific herd requirements.

CLASS/ STAGE OF PRODUCTION	ACTION/VACCINATION	ELANCO PRODUCT OPTIONS	DATE OF ADMINISTRATION AND COMMENTS
Cows: Pre-breeding	Respiratory/ Reproductive Vaccine	Inactivated: Vira Shield® 6 VL5 HB OR Vira Shield 6 L5 HB Other Vira Shield combinations are available. Consult your herd health veterinarian.	
	Parasite/Fly Control	Consult your herd veterinarian regarding prudent and effective deworming strategies for your individual operation(s). Co-Ral® Fly & Tick Spray OR CyLence® OR Clean Up™ II Pour-On	
		Apply two Corathon® OR CyLence Ultra insecticide ear tags.	
		Start feeding Rabon® Oral Larvacide in free choice mineral 3-4 weeks before fly season.	
Females: Pregnancy Check	Scours Vaccine	Scour Bos® 9	
	Respiratory/ Reproductive vaccines	Inactivated: Vira Shield 6 L5 HB Other Vira Shield combinations are available. Consult your herd health veterinarian.	
	Parasite/Fly Control	Cysectin Pour-On Clean Up II Pour-On with follow-up 2-3 weeks later for optimal louse control OR Cysectin Injectable Clean Up II Pour-On Insecticide with IGR	
Bulls: Pre-Turnout	Bull Breeding Soundness Examination, Trich Test	Contact your herd health veterinarian about conducting an exam prior to turn out.	
	Respiratory/ Reproductive vaccines	Inactivated: Vira shield 6 VL5 HB OR Vira Shield 6 L5 HB Other Vira Shield combinations are available consult your herd health veterinarian.	
	Foot Rot	Fusogard®	
	Parasite/Fly Control	Cysectin Pour-On OR Cysectin Injectable Permethrin CDS Pour-On OR CyLence OR Clean Up II Pour-On	
Apply two Corathon OR CyLence Ultra insecticide ear tags.			
Start feeding Rabon Oral Larvacide in free choice mineral 3-4 weeks before fly season.			
If Needed	Pink Eye	Pinkeye Shield® XT4	
	<i>Histophilus somni</i>	Somnu Shield®	

For all products: The label contains complete use information, including cautions and warnings. Always read, understand and follow the label and use directions.

Directions for Use: Implants are indicated for increased rate of weight gain; see product labels for full indications. Administer one dose in the ear subcutaneously according to label directions.

See label for revaccination interval recommendations.

* If you dewormed in the spring you may not need to deworm in the fall. Talk to your herd health veterinarian regarding a strategic deworming program for your herd.

Labels contain complete use information, including cautions and warnings. Always read, understand and follow the label and use directions.

** Dependent on region.

All product names, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates. ©2021 Elanco. PM-US-21-1400(2)