



PRO-BAC-C™ (MEDICATED)

Use **PRO-BAC-C MEDICATED** in dairy calves to help improve feed efficiency and increase rate of weight gain.

Controls coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in Dairy Calves.

ACTIVE DRUG INGREDIENTS:

Lasalocid (as lasalocid sodium) 6,400 grams/ton

GUARANTEED ANALYSIS:

Vitamin A, Min. 2,800,000 IU/lb
Vitamin D3, Min. 1,640,000 IU/lb
Vitamin E, Min. 988 IU/lb
Vitamin B12 Min. 8,250 mcg/lb
Thiamine, Min. 823 mg/lb
Ascorbic Acid (vit. C), Min. 3,200 mg/lb
Niacin, Min. 4,940 mg/lb
Folic Acid, Min. 204 mg/lb
Choline, Min. 4,116 mg/lb

Live Microbials

Lactic Acid Bacteria, Min. 1.35 billion CFU/lb
(*L. Acidophilus*, *L. Casei*, *E. Faecium*, *B. Thermophilum*)
Yeast, Min. (*saccharomyces cerevisiae*) 110 billion CFU/lb

PRO-BAC-C MEDICATED contains a source of live (viable) naturally occurring microorganisms.

WARNING: A withdrawal period has not been established for this product in pre-ruminating cattle.
Do not use in calves to be processed for veal.

CAUTION:

Store in a cool, dry place. For veterinary animal use only. Keep out of reach of children.
Do not give to horses.

INGREDIENTS:

Dextrose, active dry yeast (*saccharomyces cerevisiae*), vitamin A acetate, niacinamide, sodium aluminosilicate, choline bitartrate, ascorbic acid, vitamin D3 supplement, dl-alpha tocopheryl acetate (source of vitamin E), thiamine mononitrate, vitamin B12 supplement, dried *aspergillus oryzae* fermentation product, folic acid, dried *lactobacillus acidophilus* fermentation product, dried *lactobacillus casei* fermentation product, dried *bifidobacterium thermophilum* fermentation product and dried *enterococcus faecium* fermentation product.

FEEDING DIRECTIONS:

(Small measure enclosed approximates 1/8 oz).

(Cup measure enclosed approximates 6 oz).

DAIRY CALVES:

Add **PRO-BAC-C MEDICATED** to the milk replacer liquid at the rate of 1/8 oz twice daily for a 110 lb (50 kg) calf to provide 50 mg of lasalocid. Increase or decrease proportionately based on calf size to provide 1 mg lasalocid per kg of calf body weight. **Do not use in combination with other sources of lasalocid.**

CAUTION:

The safety of lasalocid in unapproved species has not been established; do not allow horses or other equines access to lasalocid as ingestion may be fatal. Feeding undiluted, mixing errors or inadequate mixing (recirculation or agitation of liquid supplements) resulting in excessive concentration of lasalocid could be fatal.

Manufactured for:

MED-VET PHARMACEUTICALS, Ltd.

14101 W. 62nd St. • Eden Prairie, MN 55346

1-833-809-4848 • www.medvetpharm.com

(Rev. 121422)