



PRO-BAC-C MEDICATED

Use **PRO-BAC-C MEDICATED** in dairy calves to help improve feed efficiency and increase rate of weight gain. Controls coccidiosis caused by Eimeria bovis and Eimeria zuernii in Dairy Calves.

ACTIVE DRUG INGREDIENTS:

Lasalocid (as lasalocid sodium) 6,400 grams/ton

GUARANTEED ANALYSIS (PER LB)

Vitamin A, Min.....	2,800,000 IU
Vitamin D3, Min.....	1,640,000 IU
Vitamin E, Min.....	988 IU
Vitamin B12 Min.....	8,250 mcg
Thiamine, Min.....	823 mg
Ascorbic Acid (Vit. C), Min.....	3,200 mg
Niacin, Min.....	4,940 mg
Folic Acid, Min.....	204 mg
Choline, Min.....	4,116 mg

Live Microbials (PER LB)

Lactic Acid Bacteria.....	1.35 billion CFUs
<small>(L. Acidophilus, L. Casei, E. Faecium, B. Thermophilum)</small>	
Yeast.....	110 billion CFUs
<small>(Saccharomyces Cerevisiae)</small>	
<small>CFUs (Colony Forming Units)</small>	

PRO-BAC-C MEDICATED contains a source of live (viable) naturally occurring microorganisms.

WARNING: A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

CAUTION:

Store in a cool, dry place.
For veterinary animal use only.
Keep out of reach of children.
Do not give to horses.

INGREDIENTS: Vitamin A acetate, cholecalciferol (source of vitamin D3 activity), dl-alpha tocopheryl acetate (source of vitamin E), vitamin B12 supplement, niacin supplement, choline chloride, folic acid, thiamine mononitrate, ascorbic acid, dextrose, dried saccharomyces cerevisiae fermentation product, dried lactobacillus acidophilus fermentation product, dried lactobacillus casei fermentation product, dried enterococcus faecium fermentation product, dried bifidobacterium thermophilum fermentation product, dried aspergillus oryzae fermentation extract and sodium silico aluminate.

FEEDING DIRECTIONS:

(Small measure enclosed approximates 1/8 oz).
(Cup measure enclosed approximates 6 oz).

DAIRY CALVES:

Add **PRO-BAC-C MEDICATED** to the milk replacer liquid at the rate of 1/8 oz twice daily for a 110 lb (50 kg) calf to provide 50 mg of lasalocid. Increase or decrease proportionately based on calf size to provide 1 mg lasalocid per kg of calf body weight. **Do not use in combination with other sources of lasalocid.**

CAUTION: The safety of lasalocid in unapproved species has not been established; do not allow horses or other equines access to lasalocid as ingestion may be fatal. Feeding undiluted, mixing errors or inadequate mixing (recirculation or agitation of liquid supplements) resulting in excessive concentration of lasalocid could be fatal.

Manufactured for:

MED-VET PHARMACEUTICALS, Ltd., 14101 W. 62nd St., Eden Prairie, MN 55346

1-800-366-8986

www.medvetpharm.com

(Rev. 102920)