

GlycoFlex® 3 Canine Formula Joint Support

VetriScience®

Ultimate Strength Stage 3 Veterinarian Formulated

A GlycOmega™ Perna Product For Joint and Connective Support

- Stage 3 Ultimate Strength
- Joint Support
- Dogs
- All Weight Ranges
- 120 chewable Tablets

Product Details:

GlycoFlex® 3 represents Stage 3 of our comprehensive stage of life program for dogs needing advanced joint support. These delicious natural imitation roast chicken flavored chewable tablets are recommended by veterinarians for dogs with a maximum need for joint support, geriatric and working dogs as well as to support normal recovery after orthopedic surgery.

Product Facts

Active Ingredients Per Tablet (4.8 g):

Glucosamine HCl (Shrimp and Crab)	1000 mg
Methylsulfonylmethane (MSM)	1000 mg
<i>Perna canaliculus</i> (GlycOmega™ brand Green-Lipped Mussel).....	600 mg
N,N-Dimethylglycine HCl (DMG)	100 mg
dl-alpha Tocopheryl Acetate (Vitamin E)	50 IU
Calcium Ascorbate (Vitamin C)	30 mg
Manganese (as Manganese Amino Acid Chelate)	10 mg
Grape Seed Extract	5 mg
L-Glutathione	2 mg
Selenium (as Sodium Selenite)	0.002 mg

Inactive Ingredients: brewers yeast, dicalcium phosphate, microcrystalline cellulose, natural imitation roast chicken flavor, silicon dioxide, stearic acid.

GlycOmega™ is a Trademark of Aroma New Zealand LTD.

For use in dogs only

Directions For Use:

Initial: (4-6 Weeks)

Up to 30 lbs:	1/2 tablet daily
31-60 lbs:	1 tablet daily
61-100 lbs:	2 tablets daily
101 lbs and over:	2 ½ tablets daily

Maintenance: (After initial period)

Up to 30 lbs:	½ tablet every other day
31-60 lbs:	1/2 tablet daily
61-100 lbs:	1 tablet daily
101 lbs and over:	1 ½ tablets daily

If giving more than 1 tablet, divide between AM and PM.

Warnings:

Not for human consumption. Keep out of reach of children and animals. In case of accidental overdose, contact a health professional immediately.

Cautions:

If animal's condition worsens or does not improve, stop product administration and consult your veterinarian. Safe use in pregnant animals or animals intended for breeding has not been proven. Administer during or after the animal has eaten to reduce incidence of gastrointestinal upset.

Store in a cool, dry place.