

# Quick Clean Disinfectant

## Zep Commercial®

Kills 99.9% of Bacteria in 5 Seconds\*\*

\*\*Staphylococcus aureus (Staph), Klebsiella pneumoniae

Cleans, Disinfects, Deodorizes

- Cuts Through Tough Grime
- Kills Cold† and Flu Viruses

†Rhinovirus

### ACTIVE INGREDIENTS

Alkyl (60% C<sub>14</sub>, 30% C<sub>16</sub>, 5% C<sub>12</sub>, 5% C<sub>18</sub>) dimethyl benzyl ammonium chlorides

..... 0.15%

Alkyl (68% C<sub>12</sub>, 32% C<sub>14</sub>) dimethyl ethyl benzyl ammonium chlorides ..... 0.15%

**INERT INGREDIENTS** ..... 99.70%

Total ..... 100.00%

KEEP OUT OF REACH OF CHILDREN.

### CAUTION

### ZEP COMMERCIAL® QUICK CLEAN DISINFECTANT

The smart way to clean and disinfect. Cleans, disinfects and deodorizes. Cuts through tough grime.

**This product is intended for use in:** Households, homes residences, ††Household kitchens, ††Institutional kitchens, Institutional bathrooms, nursery, day care centers, grocery stores, schools, health clubs, gyms, athletic facilities, barber shops, veterinary clinics; animal hospitals, kennels, <sup>1</sup>Dental Office, <sup>1</sup>Doctor's Office, home health care, animal life science laboratories, breeding establishments, grooming establishments, pet animal quarters, pet shops.

**Use on hard, non-porous surfaces:** Stainless steel, Formica® glass, glazed porcelain, plastic, sealed granite, sealed stone. This product is formulated to disinfect hard, non-porous, inanimate environmental surfaces such as floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic tile, plastic surfaces, bathrooms, shower stalls, bathtubs, and cabinets.

### Do not use on unpainted wood

- Kills Carbapenem Resistant Enterobacteriaceae (CRE) +++
- Kills Rhinovirus, causative agent of the common cold in 2 minutes!
- Kills Pandemic 2009 H1N1 Influenza A virus (formerly called swine flu)
- Kills Rhinovirus, Norovirus, Canine Parvovirus in 2 minutes!

When used as directed, Zep Commercial Quick Clean Disinfectant contains 3000 ppm of active germicide making it effective against a wide variety of microorganisms, including bacteria, antibiotic resistant bacteria and viruses.

This product will not harm most surfaces: including acrylic, sealed fiberglass and vinyl. **Do not use on unpainted wood.**

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is not for use on critical and semi-critical medical device surfaces.

Fill or refill the spray bottle with this product. Pour product into spray bottle over sink or tub basin.

**GENERAL CLEANING:** Spray soiled area then wipe with a dry paper towel or lint-free cloth.

**SANITIZE NON-FOOD CONTACT SURFACES and DEODORIZE:** To sanitize/deodorize hard, nonporous surfaces, hold container six to eight inches from the surface to be treated. Spray area until it is covered with the product. **Allow surface to remain visibly wet for 5 seconds.** Zep Commercial Quick Clean Disinfectant is an effective sanitizer against *Staphylococcus aureus* (Staph) and *Klebsiella pneumonia* (ATCC 4352). Let surface dry. For heavily soiled surfaces, a pre-cleaning step is required.

**DISINFECTING, DEODORIZING:** Remove gross filth or heavy soil prior to application of Zep Commercial Quick Clean Disinfectant. Apply product with a hand pump trigger sprayer. Hold sprayer six to eight inches from the surface to be treated. Spray area until it is covered with the product. **Allow surface to remain visibly wet for 2 minutes.** No scrubbing necessary. Allow treated surfaces to air dry or wipe off with a clean cloth.

When using on food contact surfaces: counter tops, stovetops, highchairs, kitchen appliances, thoroughly rinse all treated surfaces with potable (tap) water. This product must not be used to clean eating utensils, glassware and dishes.

**Zep Commercial Quick Clean Disinfectant is an effective disinfectant against:**

• *Pseudomonas aeruginosa* • *Staphylococcus aureus* (Staph) • *Salmonella enterica* (Salmonella) • Methicillin Resistant *Staphylococcus aureus* (MRSA) • Vancomycin Resistant *Enterococcus faecium* • Multi-Drug Resistant (MDR) *Acinetobacter baumannii* Ceftazidime and Gentamicin Resistant Extended Spectrum Beta-Lactamase *Escherichia coli* • *Klebsiella pneumoniae* New Delhi Metallo-Beta Lactamase (NDM-1) Carbapenem Resistant • Kills Carbapenem Resistant Enterobacteriaceae (CRE) +++

**Animal Housing Facilities and Poultry and Swine Premises:** Zep Commercial Quick Clean Disinfectant removes dirt, grime, blood, urine, fecal matter and other common soils found in animal housing facilities, grooming facilities, kennels, pet stores, veterinary clinics, laboratories or other small animal facilities. It also eliminates odors leaving surfaces smelling clean and fresh. It cleans, disinfects and deodorizes hard, non-porous inanimate surfaces: plated or stainless steel, aluminum, chrome, glazed porcelain, glazed tile, laminated surfaces associated with floors, walls, countertops, cages, crates, kennels, and animal equipment found in animal housing facilities.

**Disinfection of animal quarters and kennels:** To disinfect pre-cleaned animal quarters and kennels, apply Zep Commercial Quick Clean Disinfectant. Remove all animals and feed from premises. Remove all litter and droppings from floors,

walls and surfaces of facilities occupied or traversed by animals. Empty all troughs, racks, and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. **Saturate the surfaces with the disinfecting solution for a period of 2 minutes** (for Canine Parvovirus). Ventilate building and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

**\*VIRUCIDAL ACTIVITY:** This product is an effective virucide on hard non-porous surfaces against Human Immunodeficiency Virus Type-1 (HIV-1) and Avian Influenza Virus (H5N1) NIBRG-14, when the treated surface is allowed to remain wet for **30 seconds**. This product is an effective virucide on hard non-porous surfaces against Human Rotavirus (ATCC VR-2018), Hepatitis B Virus (HBV), Hepatitis C Virus (HCV), Duck Hepatitis B Virus (DHBV) and Bovine Viral Diarrhea Virus (BVDV) when the treated surface is allowed to remain wet for **1 minute**. This product is an effective virucide on hard non-porous surfaces against Feline Calicivirus, Norovirus (Norwalk Virus), Canine Parvovirus, Rhinovirus Type 14 and Rhinovirus Type 39, when the treated surface is allowed to remain to wet for **2 minutes**. Allow treated surfaces to air dry or wipe off with a clean cloth.

**KILLS HIV-1, HBV AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS**

in health care settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1, Hepatitis B Virus (HBV) and Hepatitis C Virus (HCV).

**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV AND HCV OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.**

**PERSONAL PROTECTION:** When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks, and eye coverings.

**CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of the product.

**CONTACT TIME:** Allow surfaces to remain wet for 2 minutes. (HIV-1 is inactivated in 30 seconds, HBV and HCV are inactivated in 1 minute)

**DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

**EFFICACY TESTS HAVE DEMONSTRATED THAT THIS PRODUCT IS AN EFFECTIVE BACTERICIDE AND VIRUCIDE IN THE PRESENCE OF ORGANIC SOIL (5% BLOOD SERUM).**

**For Ultrasound Transducers and Probes:**

Use only on external surfaces of ultrasound transducers and probes. With a clean washcloth, clean gel from external surface of the ultrasound transducer or

probe to be disinfected. Then thoroughly wet probe with spray. Treated surface must remain wet for 2 minutes. Let air dry.

### **PRECAUTIONARY STATEMENTS**

Hazards to Humans and Domestic Animals

**CAUTION:** Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

### **FIRST AID**

Have the product container or label with you when calling poison control center or doctor, or going for treatment.

**IF IN EYES:** Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses if present, after the first 5 minutes, then continue rinsing. Call poison control center or doctor for treatment advice. **IF ON SKIN:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

### **STORAGE AND DISPOSAL**

Do Not Contaminate Water, Food, Or Feed By Storage Or Disposal.

**PESTICIDE STORAGE** – Store in a dry place inaccessible to children.

**CONTAINER DISPOSAL** – Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or puncture and dispose in a sanitary landfill, incineration, or burning, if allowed by state and local authorities. If burned, stay out of smoke.

**PESTICIDE DISPOSAL** – Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of be use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL FOR REFILL SPRAY BOTTLE** – Do not reuse empty spray bottle except with refill container, otherwise wrap spray bottle and discard in trash. Offer for recycling if available.

+++Klebsiella pneumoniae New Delhi Metallo-Beta Lactamase (NDM-1)  
Carbapenem Resistant.

# When using on food contact surfaces: thoroughly rinse all treated surfaces with potable (tap) water.