

# Sentry<sup>®</sup> FiproGuard<sup>®</sup> Plus For Dogs (23 to 44 lbs)

## Sergeant's

For use ONLY on dogs 23 to 44 lbs

6 Applications

**Compare to Frontline<sup>®</sup> Plus for dogs, contains the same active ingredients\***

**Kills fleas, flea eggs, flea larvae, ticks & chewing lice**

- Fast acting
- Long lasting
- Waterproof

### ACTIVE INGREDIENTS:

Fipronil	9.80%
(S)-methoprene	8.80%
<b>OTHER INGREDIENT:</b>	81.40%
<b>TOTAL</b>	<b>100.00%</b>

\* This product is not manufactured by or distributed by Merial, the maker of Frontline<sup>®</sup> Plus for dogs. Frontline<sup>®</sup> is a registered trademark of Merial.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**DO NOT USE ON CATS.**

**READ ENTIRE LABEL BEFORE EACH USE. USE ONLY ON DOGS AND PUPPIES OVER 8 WEEKS OF AGE.**

### PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS. CAUTION.** Harmful if swallowed. Causes eye irritation. Avoid contact with skin, eyes, or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

**HAZARDS TO DOMESTIC ANIMALS.** For external use on dogs only. Individual sensitivities, while rare, may occur after using any pesticide product on dogs. Certain medications can interact with pesticides. Do not treat your dog with more than one pesticide product at a time. Over dosing your dog can result in serious illness and even death. Consult a veterinarian before using on medicated, debilitated, aged dogs, pregnant, or nursing dogs. Consult a veterinarian before using on dogs with known organ dysfunction.

### FIRST AID

If swallowed:	Call a poison control center or a doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor.
---------------	---

	Do not give anything by mouth to an unconscious person.
If in eyes:	Flush eyes with plenty of water. Call a doctor if irritation persists.
If on skin:	Wash with plenty of soap and water. Get medical attention if irritation persists.

## HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-781-4738 for emergency medical treatment information.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT ALLOW CHILDREN TO APPLY PRODUCT. TO PREVENT HARM TO YOU AND YOUR DOG, READ ENTIRE LABEL BEFORE EACH USE. FOLLOW ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS CAREFULLY.** Do not use on cats, rabbits or animals other than dogs. Do not allow your dog to ingest this product. Do not use on puppies under 8 weeks of age or less than 23 lbs. Use entire contents of tube on each dog. Do not split one tube between dogs. Do not use multiple tubes on one dog. Weigh your dog to be sure you are applying the right dose formulated for the weight of your dog. Separate the treated dog from all other dogs and cats for 24 hours after treatment has been applied.

Side Effects: Monitor your dog after application. Some dogs may experience temporary startle effects when any product is applied. Dogs may experience some temporary irritation at the site of product application such as redness, scratching or other signs of discomfort. If signs of sensitivity occur, bathe your dog with a mild soap and rinse with large amounts of water. If these or other side effects occur, consult your veterinarian or call 1-800-781-4738. If your dog has an unusual reaction to the initial application, consult your veterinarian before repeating application. The most common signs of incidental ingestion are excessive salivation and foaming at the mouth. If these symptoms occur, immediately feed your dog and continue to monitor your dog for the next 24 hours.

**DO NOT USE ON CATS.** Keep cats away from treated dogs for 24 hours. If applied to a cat or ingested by a cat, contact your veterinarian.

**How to open:** Remove product tubes from the package. Separate one tube from the others. Holding the tube with notched end pointing up and away from the face and body, cut off the narrow end at the notches with scissors.

**How to apply:** Invert tube over dog and use open end to part dog's hair. Squeeze tube firmly to apply all of the solution to the dog's skin, as directed below. Do not apply more often than once every 4 weeks.

**Only for Dogs Weighing 23 to 44 lbs.** Apply one tube (1.34 mL) (0.045 fl oz) as a spot on the dog's back between the shoulder blades.



**FREQUENCY OF APPLICATION**

Use this product monthly for complete control of flea, tick, and chewing lice infestations. Studies show that this product kills adult fleas for up to three months. Apply monthly if your dog is at high risk for flea reinfestation. Apply monthly to control ticks and chewing lice. This product remains effective even after bathing, water immersion, or exposure to sunlight. Allow treated area to dry thoroughly. Do not reapply for 30 days.