

# Farm & Dairy CV-90D Pest Control Spray

## Country Vet®

### Controls pests for up to 4 weeks

**Kills:** Houseflies, Fruit Flies, Mosquitos, gnats, Black Flies, Lady Bugs, and Small Flying Moths.

**Kills:** Cockroaches, Waterbugs, Palmetto Bugs, Silverfish, Firebrats, Crickets, Scorpions, Spiders, Centipedes, Sowbugs and Pillbugs.

### Active Ingredient:

Permethrin\*.....0.50%

**Other Ingredients\*\***.....99.50%

100.00%

\*Cis/trans ratio; Max 55% (+/-) and Min. 45% (+/-) trans.

\*\* Contains petroleum distillate.

Pramex® is a registered trademark of McLaughlin Gormley King Company.

**(To the User: If you can not read English, do not use this product until the label has been fully explained to you.)**

### Keep Out of Reach of Children

**Ants:** Spray trails, nests and points of entry. Spray on ants where possible. Re-treat if infestation reoccurs, but not more than once every two weeks.

**Asian Lady Birsd Beetles (Lady Bugs):** Spray insects directly around doors, windows and areas where insects can enter home.

**Indian Meal Moths, Angoumois Grain Moths:** Close doors and windows, direct spray into all parts of room especially around stored food product containers, pallets, and darkened areas. Spray 7-10 seconds for each 1100 cubic feet of space. Keep area closed for 15 minutes flowing treatment. Spray to the point of dampness but not runoff. Re-treat if infestation occurs, but not more than once every two weeks.

EXPOSED STAGES OF **Confused Flour Beetles, Saw-Toothed Grain Beetles, Red Flour Beetles, Rice Weevils, Granary Weevils, Grain Mites:** Spray into cracks and crevices, around cartons or containers or other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Spray to the point of dampness but not run off. Re-treat if infestation occurs, but not more than once every two weeks. Infested stored food products in the home should be disposed of.

**Carpet Beetles:** Spray directly on beetles, floors, baseboards, shelves, and carpets. Spray to the point of dampness but not runoff. Re-treat if infestation occurs but not more than once every two weeks.

**Fleas & Ticks (Adult and Larve):** Remove and thoroughly clean or destroy pet bedding. Spray sleeping quarters, bed, entire floor and floor covering, with special attention to cracks and crevices around baseboards and window and door frames. Infested upholstered furniture may also be treated with this product. Spray to the point of dampness but not runoff. A test application should be made to upholstery or drapery fabrics in an inconspicuous place before use.

To kill **Lice and Louse Eggs:** Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray each square foot for three seconds. Allow all sprayed articles to dry thoroughly before use. \*Spray only on those garments and parts of bedding, including mattresses and furniture that can not either be laundered or dry cleaned. Do not use on sheets or pillow cases.

**For Direct Application to Dogs (Over 12 Weeks of Age):**  
**TO CONTROL Fleas, ticks, and Lice.**

Hold the container upright. Spray at a distance of 1-12 inches. Starting at the tail, spray over entire haircoat while moving the dispenser rapidly; making sure that the coat is thoroughly wetted to the skin. While spraying, fluff the hair so the spray will penetrate to the skin. Make sure spray wets ticks thoroughly. Spray against the lay of the hair, roughing the hair on long-haired dogs. Do not spray into the eyes, face or genitalia. For best results in killing ticks, spray directly on the tick. Allow dog to air dry or blow dry. Do not reapply more than once every 14 days.

Use protective gloves or mitts when applying product. **DO NOT USE ON CATS OR KITTENS.**

**In Animal Quarters:** For rapid control of **Houseflies, Mosquitos, and Small Flying Moths**, close windows and doors and spray at the rate of 2 seconds per 100 cubic feet of space. For best results, keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is completed. Do not contaminate animal feed or water during application.

**To Treat Livestock:** To protect beef cattle, dairy cattle, goats sheep, hogs and horses from attacks of **Stable Flies, Horse Flies, Deer Flies, Face Flies, House Flies, Horn Flies, Mosquitos and Gnats:** Spray about 3 seconds to each side being careful to spray back, withers, and forelegs thoroughly. To protect from **Face Flies**, spray the face and head, but do not spray into the eyes. Repeat treatment when flies are troublesome.

To Kill **Screw Worms** infesting Beef cattle, horses and sheep, hold can about 4-8 inches from area to be treated with nozzle pointed at wound. Spray the area until it is completely wet- three to ten seconds are usually enough. The area surrounding the wound should also be sprayed. This product is not recommended for use in deep wounds. Some strains of screw worms may develop resistance to pyrethroids. This will be observed by lack of response to treatment. Where pyrethroid resistance is known or suspected, this product should not be used.

FOR SWINE: Do not ship animals for slaughter within 5 days of last treatment.

TO CONTROL **Blood Suckling Lice**: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every three weeks if required.

TO CONTROL **Poultry Lice**: Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist. For control of **Bed Bugs and Mites** in Poultry houses, spray crevices of roost poles, cracks in walls and cracks in nests where the bed bugs and mites hide. Spray to the point of dampness but not runoff.

FOR THE CONTROL OF GARDEN INSECTS: For use on roses, dahlias, asters, and other ornamentals to kill **Japanese Beetles** hit by spray, as well as **White Flies, Aphids, Exposed Thrips, Spider Mites, Red Mites, and Leafminers**.

May be used on plants, as well as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camelias, carnations, chrysanthemums, dahlias, delphiniums, dogwood, English ivy, euonymus, fuchsia, geraniums, crassula, grape vine, Kentia palm, laurel, marigolds, rhododendrons, roses, rubber plants, snap dragons, stocks, wandering Jews and zinnias.

Effective when used as directed against **White Flies, Aphids, Spider and Red Mites, Exposed Thrips, Lace Bugs, Japanese Beetles, Pavement Ants, Clover Mites and Armyworms**. When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Spray in 1 second bursts. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F (32°C). Over application may cause damage. Contact insects directly whenever possible. Re-treat if infestation occurs, but not more than once every two weeks.

### **STORAGE AND DISPOSAL**

**Pesticide Storage:** Store in a cool, dry area away from children and pets and away from heat, sparks and open flame. Do not transport or store below 32°F (0°). Do not freeze.

**Pesticide disposal and container handling:** Do not puncture or incinerate. **If empty:** place in trash or offer for recycling if available. **If partially filled:** Call your local solid waste agency for disposal instructions.

## **FIRST AID**

### **If Swallowed**

- Immediately call a Poison Control Center or doctor
- Do not induce vomiting unless told to do so by a poison control center or doctor
- Do not give **any** liquid to the person
- Do not give anything by mouth to an unconscious person

### **If in Eyes**

- Hold eye open and rinse slowly and gently with water for 15-20 minutes
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye
- Call a poison control center or doctor for treatment advice

### **If on Skin or Clothing**

- Take off contaminated clothing
- Rinse skin immediately with plenty of water for 15-20 minutes
- Call a poison control center or doctor for treatment advice

## **HOTLINE NUMBER**

Have the product or label with you when calling a poison control center or doctor, or going for medical treatment. For additional information on this product (including health concerns, medical emergencies or pesticide incidents), you may call 1-888-740-8712.

## **NOTE TO PHYSICIAN**

Contains petroleum distillate- vomiting may cause aspiration pneumonia.

## **PRECAUTIONARY STATEMENTS**

### **Hazardous to Humans & Domestic Animals**

#### **Caution**

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

## **Environmental Hazards**

This product is extremely toxic to aquatic organisms, including fish and invertebrates. Do not apply directly to or near water. Drift and runoff may be hazardous to fish in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater, or rinsate. See directions for use for additional precautions and requirements.

This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift blooming crops or weeds while bees are actively visiting the treatment area.

## **PHYSICAL OR CHEMICAL HAZARDS**

Contents under pressure. Do not store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F (54°C). may cause bursting.

## **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with it's labeling.

## **READ ENTIRE LABEL BEFORE EACH USE**

### **Use Restrictions**

- Except when applying to livestock, do not allow adults, children or pets to enter until vapors, mists and aerosols have dispersed and the treated area has been thoroughly vented.
- Except when applying to livestock, do not allow adults, children or pets to enter the treated area until sprays have dried.
- In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.
- Remove pets, birds and cover any water inhabited by fish (such as aquariums and ornamental fish ponds) before application. Turn off aquarium filtering systems before spraying and wait approximately 15 minutes after application to turn back on.
- **This product is not intended for use on humans.** If lice infestations should occur on humans, use a product labeled for use on humans.
- Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter ant drain during or after application.

## **SHAKE WELL BEFORE USING**

Remove protective cap, hold container upright with nozzle away from you. Depress valve and spray from a distance of 8-10 inches (12-15 inches for baseboards, moldings & floors). Test product on an inconspicuous area before applying to check for staining or discoloration. Spray until surface is slightly wet, but not to the point of runoff. Over application may cause damage. Will not stain surfaces or fabric where water alone causes no damage.

### **Houseflies, Fruit Flies, Mosquitos, Gnats, Black Flies, Skipper Flies, Blow Flies, and Small Flying Moths:**

Close all doors and windows and spray room at the rate of 2-3 seconds per 100 cubic feet. Direct spray to all parts of the room, especially windows and other light sources, which attract these insects. Keep nozzle in constant motion. Re-treat if infestation occurs, but not more than once every two weeks.

**Cockroaches, Waterbugs, Palmetto Bugs, Silverfish, Firebrats, Crickets, Scorpions, Spiders, Millipeds, Centipedes, Sowbugs and Pill Bugs:** Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Spray to the point of dampness but not runoff. Re-treat if infestation occurs, but not more than once every two weeks.